

# Ectospec

Cypermethrin Technical 2.5%w/v

## INDICATIONS

For the control of lice and flies on cattle.

## BENEFITS

- ✓ Ready to use insecticide for cattle
- ✓ For fly and lice control in cattle
- ✓ No milk withdrawal period
- ✓ 10 days meat withdrawal
- ✓ Low dose rate 10ml/treatment



## PACKAGING

List No	Unit Package	Case Size
1ECT001	500ml	12
1ECT002	2.5L	4



# Ectospec

Cypermethrin Technical 2.5%w/v



## PRESENTATION

A clear, pale yellow, oily solution containing a 2.5% w/v Cypermethrin technical (93%) (cis:trans 50:50) for topical administration as a pour on application.

## USES

For the control of lice and flies on cattle.

## DOSAGE & ADMINISTRATION

For topical administration as a pour on.

### Fly Control

Apply a single 10ml dose at the onset of the fly season and repeat as necessary at 5 to 8 week intervals

Frequency of administration may, however, have to be varied depending on the level and type of infestation.

The 10ml dose should be applied at an even rate along the backline from the crown of the head to the top of the rump.

### Lice Control

A single 10ml dose is normally sufficient to control lice. A few lice may survive on a small minority of animals. In cases of heavy challenge, if necessary, a repeat dose may be applied after 4 weeks. The 10ml dose should be applied at an even rate along the backline from the crown of the head to the top of the rump.

## UNDESIRABLE EFFECTS (GENERAL)

Occasionally, slight signs of discomfort may be observed in some cattle during the 48 hours following application. These side effects are only temporary and have no long-term implications.

## USE DURING PREGNANCY AND LACTATION

The active has a low oral toxicity and very little transdermal absorption is expected, and to date there has been no reported complications in pregnancy or lactation.

## INTERACTION WITH OTHER MEDICAMENTS AND OTHER FORMS OF INTERACTIONS

It is not anticipated that Ectospec will interact with any of the medicines commonly administered to cattle.

## PROTECTION OF OPERATORS

Ectospec should be administered in a well ventilated area. Protective clothing, including rubber gloves and rubber boots, should be worn and any accidental splashes on skin or eyes should be washed off immediately. Do not eat, drink or smoke when applying Ectospec. Wash hands and any exposed skin before eating, drinking or smoking and immediately after application has finished.

## SPECIAL PRECAUTIONS FOR USE

Dangerous to fish and other aquatic animals. Do not contaminate ponds, waterways or ditches with the product or used containers. Unused product or containers should be disposed of in accordance with current practice for pharmaceutical waste under national waste disposal regulations.

## OVERDOSE

Care should be taken not to overdose. Overdosing may invalidate the stated milk and meat withholding times.

## WITHDRAWAL PERIODS

Animals must not be slaughtered for human consumption during treatment. Treated animals should not be slaughtered for human consumption for a period of 10 days after treatment.

No withdrawal time is necessary for milk, although cows should be treated immediately after milking to allow as long a time as possible to elapse before the next milking.

## PHARMACEUTICAL PRECAUTIONS

Incompatible with alkali materials  
Do not store above 25°C.  
Keep out of reach and sight of children.

## LEGAL CATEGORY

LM

## PACKAGE QUANTITIES

500ml multidose and 2.5L containers.

## MARKETING AUTHORISATION NUMBER:

VPA 10960/79/1

## MARKETING AUTHORISATION HOLDER:

Cross Vetpharm Group Ltd.