

EQUIBACTIN

Trimethoprim 66.7mg/g
Sulfadiazine 333.3mg/g

INDICATIONS

Treatment of infections in horses caused by bacteria sensitive to the combination of trimethoprim and sulfadiazine, particularly:
 Respiratory tract infections associated with *Streptococcus* spp. and *Staphylococcus aureus*;
 Gastrointestinal infections associated with *E. coli*;
 Urogenital infections associated with beta-hemolytic streptococci;
 Wound infections and open or drained abscesses associated with *Streptococcus* spp. and *Staphylococcus aureus*.

BENEFITS

- Palatable aniseed flavour
- Single syringe pack, easy to administer
- Treats a 600kg horse in a single syringe
- Broad spectrum activity against respiratory, gastrointestinal, urogenital and wound infections in horses
- Powerful antibiotic formula of sulfadiazine and trimethoprim



PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1EQU005	45G SYRINGE	48

See reverse side for Administration and Dosage.



EQUIBACTIN VET. (333 mg/g + 67 mg/g) ORAL PASTE FOR HORSES.

Trimethoprim 66.7mg/g Sulfadiazine 333.3mg/g

PRESENTATION

Oral paste in a white to almost white suspension, containing trimethoprim 66.7mg and sulfadiazine 333.3mg per gram.

TARGET SPECIES

Horses.

INDICATIONS FOR USE, SPECIFYING THE TARGET SPECIES

Treatment of infections in horses caused by bacteria sensitive to the combination of trimethoprim and sulfadiazine, particularly:

Respiratory tract infections associated with *Streptococcus* spp. and *Staphylococcus aureus*;

Gastrointestinal infections associated with *E. coli*;

Urogenital infections associated with beta-hemolytic streptococci;

Wound infections and open or drained abscesses associated with *Streptococcus* spp. and *Staphylococcus aureus*.

CONTRAINDICATIONS

Do not use in horses known to be hypersensitive to sulfonamides, with serious hepatic or renal insufficiency nor with blood dyscrasias,

Do not use this product to treat abscesses without proper drainage.

SPECIAL PRECAUTIONS FOR USE

SPECIAL PRECAUTIONS FOR USE IN ANIMALS

To avoid under- or overdosing assess bodyweight and dosage as accurately as possible before dosing.

Do not use the same syringe in more than one animal.

Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies.

SPECIAL PRECAUTIONS TO BE TAKEN BY THE PERSON ADMINISTERING THE VETERINARY MEDICINAL PRODUCT TO ANIMALS

People with known hypersensitivity to sulfonamides should avoid contact with the veterinary medicinal product.

In case of reaction of hypersensitivity after exposure (such as skin rash), seek medical advice and show the package leaflet or the label to the physician. In case of severe reactions (swelling of the face, lips or eyes), seek prompt medical attention and take the package leaflet with you.

ADVERSE REACTIONS (frequency and seriousness)

Decrease or loss of appetite can occur in treated animals.

Loose faeces and diarrhea may develop during treatment with the product.

If such effects appear, discontinue treatment immediately and institute appropriate symptomatic measures.

USE DURING PREGNANCY AND LACTATION

Laboratory studies in rats and mice have shown evidence of teratogenic effects. The safety of the product has not been established during pregnancy. Use only according to the benefit/risk assessment by the responsible veterinarian.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

Potentiated sulfonamides used in conjunction with detomidine are known to be able to cause fatal arrhythmias in the horse.

AMOUNTS TO BE ADMINISTERED AND ADMINISTRATION ROUTES

Administration route: Oral use.

Posology:

5 mg trimethoprim and 25 mg sulfadiazine per kg body weight per day to a maximum of 5 days.

One syringe is intended for 600 kg bodyweight and each syringe is subdivided into 12 markings.

The equivalent of one marking is sufficient to treat 50 kg of bodyweight and the minimum body weight for treatment is 50 kg.

Directions for use

Horse weight should be accurately determined for the correct use of the paste.

The calculated dose is provided by adjusting the ring on the plunger according to the body weight of the horse.

The paste is administered orally by inserting the nozzle of the syringe through the interdental space and depositing the required amount of paste on the back of the tongue. The animal's mouth should be free of any food. Immediately after administration, elevate the head of the horse for a few seconds to ensure the dose is swallowed.

OVERDOSE (symptoms, emergency procedures, antidotes), if necessary

Not known.

WITHDRAWAL PERIODS

Meat and offal: 14 days

Not permitted for use in mares producing milk for human consumption.

PHARMACODYNAMIC PROPERTIES

Both active substances produce a sequential double blockade of bacterial synthesis of folic acid. This results in a synergistic and bactericidal action inhibiting sequential steps in the synthesis of purines, which are required for DNA synthesis. The combination has a broad action against many Gram-positive and Gram-negative bacteria such as staphylococci, streptococci and *E.coli*.

PHARMACOKINETIC PARTICULARS

After a single oral administration of 5 mg trimethoprim and 25 mg sulfadiazine per kg body weight to horses, the following parameters (mean ± sd) were observed:

	C _{max} (mg/ml)	T _{max} (hour)	T _{1/2} el (hour)
trimethoprim	2.35 ± 0.59	0.91 ± 0.32	2.74 ± 0.91
sulfadiazine	14.79 ± 3.47	1.90 ± 0.76	7.4 ± 1.8

Food intake appeared to affect the pharmacokinetic profile as both trimethoprim and sulfadiazine have been absorbed more rapidly in fasted horses.

Excretion of both actives is chiefly by the kidneys, by both glomerular filtration, and tubular secretion.

Urine concentrations of both trimethoprim and sulfadiazine are severalfold higher than blood concentrations. Neither trimethoprim nor sulfadiazine interferes with the excretion pattern of the other.

SHELF LIFE

Shelf-life of the veterinary medicinal product as packaged for sale: 2years.

Shelf-life after first opening the immediate packaging: 8 weeks.

SPECIAL PRECAUTIONS FOR STORAGE

Do not refrigerate or freeze.

NATURE AND COMPOSITION OF IMMEDIATE PACKAGING

Pre-filled multi-dose (LD) polyethylene syringe with adjustable screw ring closed with a (LD) polyethylene cap, packed in a cardboard box. Each syringe contains 45 g paste.

Registered with HSE

LEGAL CATEGORY

POM-VPS

VPA 10816/004/001

