

# TYLOVET 20%

*Tylosin*



## INDICATIONS

For the treatment of infections caused by microorganisms susceptible to tylosin.

## BENEFITS

- Tylosin Tartrate base, readily soluble in water
- Broad spectrum activity against Gram-positive and Gram-negative organisms in cattle, sheep and pigs
- Excellent at reducing sub clinical mastitis and SCC issues at drying off in dairy cows
- Short withdrawals, milk 4 days and meat 21 days
- A proven performer

## PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1TYL009	100ml	12

See reverse side for full indications, administration and dosage.

Bimeda<sup>®</sup>  
 Broomhill Road  
 Tallaght  
 Dublin 24, Ireland  
 Tel: 1850-515253



[www.bimeda.ie](http://www.bimeda.ie)

## TYLOVET 20% W/V SOLUTION FOR INJECTION

### PRESENTATION

A clear yellow solution for injection.  
Each ml contains:  
Tylosin (as Tylosin Tartrate) 200.0 mg

### TARGET SPECIES

Cattle, pigs.

### USES

For the treatment of infections caused by microorganisms susceptible to tylosin. The spectrum of antimicrobial activity of tylosin includes:  
Staphylococcus spp  
Streptococcus spp.  
Corynebacterium spp.  
Neisseria spp.  
Flavobacterium spp.  
Campylobacter spp.  
Bacillus anthracis  
Moraxella bovis  
Clostridium spp.  
Haemophilus spp.  
Bordetella bronchiseptica  
Spirochaetes  
Bacteroides spp.  
Mycoplasma spp.

### CONTRAINDICATIONS, WARNINGS ETC

Do not use in case of hypersensitivity to the active ingredient or to any other macrolide antibiotics.  
Do not use in new-born animals.  
Do not use in equines.  
Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.  
The safety of the veterinary

medicinal product has not been established during pregnancy and lactation.  
Use only according to the benefit/risk assessment by the responsible veterinarian.

### OPERATOR WARNINGS

Avoid skin contact with solution.

### DOSES AND ADMINISTRATION

The recommended dosage rate is 10 mg/kg bodyweight twice daily i.e. 0.5 ml per 10 kg body weight. For intramuscular administration. Treatment should be carried out for 3 to 5 consecutive days. To ensure a correct dosage, bodyweight should be determined as accurately as possible.

### OVERDOSE

Overdose should be avoided using the most accurate bodyweight estimation.

### WITHDRAWAL PERIOD(S)

Meat and offal: 21 days.  
Milk: 96 hours.

### INCOMPATIBILITIES

Do not mix with any other medicinal products.

### SHELF-LIFE

Shelf-life of the veterinary medicinal product as packaged for sale: 3 years.  
Shelf-life after first opening the immediate packaging: 14 days.

### STORAGE

Store in a refrigerator (2°C - 8°C).  
Protect from light. Do not freeze.

### PACKAGING

100 ml amber coloured type II glass vial sealed with butyl rubber stoppers and aluminium cap.

### LEGAL CATEGORY

POM

### DISPOSAL

Any unused product or waste materials should be disposed of in accordance with national requirements

VPA 10989/008/001

