
# **BOVIFERM PLUS**

DATA SHEET

**Supplementary feed** 



# **USES**

Supplementary feeding stuff for the diarrheoic calf. Boviferm Plus solution is suitable to be fed with milk. For rapid rehydration and acidosis correction.

# LIST No UNIT PACKAGE CASE SIZE 1BOV021 115g sachet 12

# **BENEFITS**

- High rehyration properties and Strong Ion Difference (SID)
- Can be mixed with milk
- Calf can remain on milk feed
- Contains Oralin, a microbial additive that stabilises the gut flora



# **BOVIFERM PLUS**



# Supplementary feed

### **PRESENTATION**

Supplementary feed for calves with diarrhoea.

For the stabilisation of water and electrolyte balance.

Compatible with milk digestion Boviferm® Plus<sup>SID</sup> provides the calf with electrolytes, buffering agents, energy, lactic acid bacteria, mucins and bulking agents as well as toxin binding substances, for rehydration.

# **COMPOSITION**

Dextrose; Whey powder; Sodium propionate, Sodium chloride; Potassium chloride; Banana flakes; Pre-gelatinised rice flour; Calcium chloride; Magnesium chloride.

#### CONTENTS

Crude protein 4.0%; Crude fibre 1.4%; Crude fat 0.5%; Crude ash 21.0%; Sodium 4.4%; Potassium 1.7%; Chlorides 3.8%; moisture 6.0%.

# **ADDITIVES PER KG.**

Bentonite Montmorillonite; Guar seed flour; Citrus pectin; Camomile; Aniseed; Fennel seed; Enterococcus faecium DSM 10663 NCIMB 10415 (E 1707) (ORALIN®) 1.75 x 1010 CFU

# **FEEDING INSTRUCTIONS**

In case of risk of, during periods of, or recovery from digestive disturbance (diarrhoea):

Twice per day: Give 1 sachet (115 gm) of Boviferm® Plus<sup>SID</sup> mixed into 2 litres of water and add 0.5 to 4 litres of milk or milk replacer (according to the calf's daily requirement).

Administer Boviferm® Plus<sup>SID</sup> until diarrhoea stops, at minimum for 2 to 3 days. Do not stop milk feeding. Additionally give water ad libitum.

It is recommended that a veterinarian's opinion be sought before use.

# **SPECIAL INSTRUCTIONS**

As Boviferm® Plus<sup>SID</sup> contains viable lactic acid bacteria, the mixing temperature should not exceed 50°C.

# **STABILITY**

24 months

# **STORAGE**

Cool and dry

# **PACK SIZE**

Box of 24 x 115 q

# **MARKETED BY**

**Bimeda Animal Health Limited** 2, 3 & 4 Airton Close, Airton Road Tallaght, Dublin 24, Ireland

Tel: +353 (0) 1 466 7900



