

ECTOSPEC 2.5% W/C

Pour-On- Solution

Cypermethrin Technical 2.5% w/v

DATA SHEET



INDICATIONS

Ectospec is a ready-to-use ecto-parasiticide for cattle which is licensed for the control of lice and flies on cattle.

BENEFITS

- Ready to use ecto-parasiticide for cattle.
- For fly and lice control.
- No milk withdrawal period.
- 10 days meat withdrawal.
- Low dose rate of 10ml/treatment.



LIST No	UNIT PACKAGE	CASE SIZE
1ECT001	500 ml	12
1ECT002	2.5L	6

See reverse for Administration & Dosage



ECTOSPEC 2.5% w/v

Pour-On Solution

Cypermethrin Technical 2.5% w/v



PRESENTATION

A clear, pale yellow, oily solution containing 2.5% w/v Cypermethrin technical (93% w/w; cis/trans isomers 50:50) as a pour on application.

USES

For the control of lice and flies on cattle.

DOSAGE & ADMINISTRATION

For topical administration as a pour on.

Fly Control

Apply a single 10ml dose at the onset of the fly season and repeat as necessary at 5 to 8 weeks intervals.

Frequency of administration may, however, have to be varied depending on the level and type of infestation.

The 10ml dose should be applied at an even rate along the backline from the crown of the head to the top of the rump.

Lice Control

A single 10ml dose is normally sufficient to control lice. A few lice may survive on a small minority of animals. In cases of heavy challenge, if necessary, a repeat dose may be applied after 4 weeks. The 10ml dose should be applied at an even rate along the backline from the crown of the head to the top of the rump.

UNDESIRABLE EFFECTS (GENERAL)

Occasionally, slight signs of discomfort may be observed in some cattle during the 48 hours following application. These side effects are only temporary and have no long-term implications.

USE DURING PREGNANCY AND LACTATION

The active has a low oral toxicity and very little transdermal absorption is expected, and to date there has been no reported complications in pregnancy or lactation.

INTERACTION WITH OTHER MEDICAMENTS AND OTHER FORMS OF INTERACTIONS

It is not anticipated that Ectospec will interact with any of the medicines commonly administered to cattle.

PROTECTION OF OPERATORS

Ectospec should be administered in a well ventilated area. Protective clothing, including rubber gloves and rubber boots, should be worn and any accidental splashes on skin or eyes should be washed off immediately.

Do not eat, drink or smoke when applying Ectospec. Wash hands and any exposed skin before eating, drinking or smoking and immediately after application has finished.

CYPERMETHRIN

Cypermethrin is a SP and has a rapid paralytic action on ectoparasites, preceded by muscular excitation and convulsions. The pyrethroids have sufficient stability to have a prolonged effect when applied to the animals.

Cypermethrin is not systemically active when used as a pour on preparation. When applied in this manner, the product spreads over the surface of the animal.

SPECIAL PRECAUTIONS FOR USE

Dangerous to fish and other aquatic animals.

Do not contaminate ponds, waterways or ditches with the product or used containers.

Unused product or containers should be disposed of in accordance with current practise for pharmaceutical waste under national waste disposal regulations.

Do not treat calves under 7 days of age.

Do not use in cases of hypersensitivity to the active substance.

OVERDOSE

Care should be taken not to overdose.

Overdose may invalidate the stated milk and meat with holding times.

WITHDRAWAL PERIODS

Animals must not be slaughtered for human consumption during treatment. Treated animals should not be slaughtered for human consumption for a period of 10 days after treatment.

No withdrawal time is necessary for milk, although cows should be treated immediately after milking to allow as long a time as possible to elapse before the next milking.

PHARMACEUTICAL PRECAUTIONS

Incompatible with alkali materials.

Do not store above 25°C.

Keep out of reach and sight of children.

LEGAL CATEGORY

POM

PACKAGE QUANTITIES

500ml multidose and 2.5L containers.

MARKETING AUTHORISATION NUMBER

VPA 22033/014/001

MARKETING AUTHORISATION HOLDER

Bimeda Animal Health Ltd

2, 3 & 4 Airton Close, Airton Road

Tallaght, Dublin 24, Ireland

Tel: +353 (0) 1 466 7900

Use Medicines Responsibly.

A full product SPC is available on request from Bimeda or alternatively can be found on the HPRA website.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

Bimeda data sheet updated: December 2024

Global Excellence in Animal Health

www.bimeda.ie

 **Bimeda**